



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

24. Proposed by D. H. DAVISON, C. E., Minonk, Illinois.


For the purpose of locating the most eligible point for a county-seat, it is required to determine the centre of a county whose dimensions are as follows: Beginning at the S. W. corner, thence E. 15 miles, thence N. $33\frac{3}{4}$ miles, thence W. 6 miles to the north end of the meridian running south through the county, thence south-westerly to a point 6 miles W. from the meridian and $9\frac{1}{2}$ miles S. of the north end of said meridian, thence S 3 miles, thence W. 3 miles, and thence S. 21 miles to the place of beginning.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science in Texarkana College, Texarkana, Arkansas; and F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

Let \bar{x} , \bar{y} , be the co-ordinates of the centroid, and divide the county into three parts as in the figure, then we easily get with A as origin

$$\bar{x} = \frac{\int_0^3 \int_0^{21} x dx dy + \int_3^9 \int_0^{19\frac{1}{2} + \frac{3}{4}x} x dx dy + \int_9^{15} \int_0^{33\frac{3}{4}} x dx dy}{\int_0^3 \int_0^{21} dx dy + \int_3^9 \int_0^{19\frac{1}{2} + \frac{3}{4}x} dx dy + \int_9^{15} \int_0^{33\frac{3}{4}} dx dy} = 8\frac{15}{8}\frac{3}{8} \text{ miles.}$$

$$\bar{y} = \frac{\int_0^3 \int_0^{21} y dx dy + \int_3^9 \int_0^{19\frac{1}{2} + \frac{3}{4}x} y dx dy + \int_9^{15} \int_0^{33\frac{3}{4}} y dx dy}{\int_0^3 \int_0^{21} dx dy + \int_3^9 \int_0^{19\frac{1}{2} + \frac{3}{4}x} dx dy + \int_9^{15} \int_0^{33\frac{3}{4}} dx dy} = 15\frac{1}{8}\frac{8}{4} \text{ miles.}$$



\therefore Measure east from beginning $8\frac{15}{8}\frac{3}{8}$ miles, then north $15\frac{1}{8}\frac{8}{4}$ miles.

[The proposition that, "The point of the area of a triangle, which has the sum of its distances to all other points of the area a *minimum*, is the centre of gravity of the area," which I think holds for other figures, practically solves problem 24, No. 2. I have made out the proof for the triangle but it occupies two pages.

R. J. ADCOCK, Larchland, Illinois.]

Also solved by P. S. BERG.

A CORRECTION.—On page 246 of the MONTHLY, my remarks in the lower four lines *above* the Note, are not true and should be expunged. They were hastily made upon insufficient investigation. In prob. 20, those remarks hold almost true, but in the general problem they can not ever be true. My solution is not at all affected by those misstatements. The solution may be more easily understood by adding, that, "when Sirius rises, some point of the ecliptic is then rising, and as the Sun is always on the ecliptic the Sun must be at that point, in order to rise synchronously with Sirius.

S. H. WRIGHT.

34. Proposed by GEORGE LILLEY, Ph. D., LL. D., Park School, Portland, Oregon.

A hare is at O , and a hound at E , 40 rods east of O . They start at the same instant each running with uniform velocity. The hare runs north. The hound runs directly towards the hare and overtakes it at N , 320 rods from O . How far did the hound run?